

Specimen Instructions

Peripheral Whole Blood


Use only tubes provided inside the FoundationOne® Liquid Specimen Collection and Shipping Kit. Other tubes will not be accepted.



Instructions For Use

Accurate analysis of cell-free DNA requires proper collection technique and handling of the sample. Failure to adhere to these instructions can compromise results by diluting cell-free DNA with DNA from white blood cell lysis.

- 1** Check special tubes provided in FoundationOne Liquid kits to confirm liquid is clear and without cloudiness or crystals.
- 2** Label tubes with date of collection and two patient identifiers.
- 3** Collect two tubes of whole blood (8.5mL per tube).
 - Prevent backflow: tubes contain chemical additives and it is important to avoid backflow into patient.
 - Collect specimen by venipuncture according to CLSI H3-A6.¹
 - Fill tubes completely (8.5mL per tube).
- 4** Remove the tube from adapter and immediately **mix by gentle inversion 8 to 10 times**. Inadequate or delayed mixing may result in inaccurate test results. One inversion is a complete turn of the wrist, 180°, and back per the figure below.



- 5** Place specimen, completed requisition form (TRF) (remember to include patient's diagnosis), insurance information, available reports, and accompanying documents into the FoundationOne Liquid Specimen Collection and Shipping Kit (copies of pathology reports and/or other clinical documentation).
 - Confirm each tube is labeled with the supplied labels indicating the date of collection and two unique patient identifiers (label included in kit).
- 6** Preferably on the same day of collection, ship via FedEx priority overnight delivery at ambient temperature. Do not freeze or refrigerate blood samples.

Temperature is important. Keep at room temperature (43-99° F, 6-37° C).

DO NOT FREEZE.

LOCAL CONTACT

MALAYSIA
Roche (Malaysia) Sdn. Bhd.
Level 21, The Pinnacle, Persiaran Lagoon,
Bandar Sunway, 47500 Subang Jaya,
Selangor Darul Ehsan.

Customer Services team: 1800 806 857
For more information, contact
our APAC Customer Services team:
APAC.foundation@roche.com

Reference

1. Clinical and Laboratory Standards Institute. H3-A6, Procedures for the collection of diagnostic blood specimens by venipuncture; approved standard-sixth edition.